

EMPOWER chart audit tool

*Note: Please complete this chart **ONLY** for patients with HR-positive HER2-negative metastatic breast cancer (mBC).*

Institution:		Patient since (month/year):		
Patient demographics				
Age:	Gender (circle): Female Male	Menopausal status (circle): Pre/perimenopausal Postmenopausal	Ethnicity:	Body mass index:
Medical history:				
Current medication list:				
Stage at <u>initial</u> diagnosis:				
T <input type="checkbox"/> 0 <input type="checkbox"/> 1 <input type="checkbox"/> 2 <input type="checkbox"/> 3	M <input type="checkbox"/> M0 <input type="checkbox"/> M1	N <input type="checkbox"/> 0 <input type="checkbox"/> 1–3 <input type="checkbox"/> 4+	Official diagnosis:	
Diagnostic testing performed:				
Molecular testing performed:				
Initial treatment for metastatic breast cancer (mBC):				
Degree of patient participation in treatment decision:				
<input type="checkbox"/> High (ie, involved in all decisions, well informed) <input type="checkbox"/> Medium (ie, understands the important aspects, though not in detail) <input type="checkbox"/> Low (ie, patient depends on provider for all decisions) <input type="checkbox"/> Unknown				

Please send completed chart audits to jmcmullen@medlearninggroup.com

1st-line treatment included:

- Fulvestrant
- Abemaciclib + fulvestrant
- Palbociclib + fulvestrant
- Ribociclib + fulvestrant
- Tamoxifen
- Ribociclib + tamoxifen
- Exemestane + everolimus
- Fulvestrant + everolimus
- Tamoxifen + everolimus
- Capecitabine
- Intravenous (IV) chemotherapy +/- capecitabine
- Aromatase inhibitor (AI)
- Fulvestrant + AI
- Palbociclib + AI
- Ribociclib + AI
- Abemaciclib + AI
- Clinical trial: _____

Factors influencing treatment decision:

- Patient preference
- Mechanism of action of therapy
- Cost of therapy
- Clinical data
- Professional experience with therapy in practice
- Other: _____

Date of treatment initiation:

Duration of treatment:

Adverse events (AEs):

Management strategy for AE(s):

Additional members of healthcare team involved in treatment decisions for this patient (please mark all that apply):

- Oncologist
- Oncology nurse
- Nurse navigator
- Pharmacist
- Social worker
- Patient caregiver/family
- Other (please specify): _____

- None of the above

Method used to detect progression:

Progression of disease?

- Yes
- No

Please describe sites and symptoms associated with progression.

Second-line treatment for mBC

Degree of patient participation in treatment decision:

- High (ie, involved in all decisions, well informed)
- Medium (ie, understands the important aspects, though not in detail)
- Low (patient depends on provider for all decisions)
- Unknown

Treatment (2nd line):

- Fulvestrant
- Abemaciclib + fulvestrant
- Palbociclib + fulvestrant
- Ribociclib + fulvestrant
- AI
- Palbociclib + AI
- Ribociclib + AI
- Abemaciclib + AI
- Tamoxifen
- Ribociclib + tamoxifen
- Exemestane + everolimus
- Fulvestrant + everolimus
- Tamoxifen + everolimus
- Capecitabine
- IV chemotherapy +/- capecitabine
- Abemaciclib
- Clinical trial _____

Factors influencing decision:

- Patient preference
- Prior therapy
- Mechanism of action of therapy
- Cost
- Clinical data
- Professional experience with therapy in practice
- Other: _____

Duration of treatment:	
AE(s):	
Management of AE(s):	
Additional members of healthcare team involved in decisions on patient treatment (please mark all that apply): <ul style="list-style-type: none"> <input type="checkbox"/> Oncologist <input type="checkbox"/> Oncology nurse <input type="checkbox"/> Pharmacist <input type="checkbox"/> Social worker <input type="checkbox"/> Patient caregiver/family <input type="checkbox"/> Other (please specify): _____ <input type="checkbox"/> None of the above 	
Progression of disease? <ul style="list-style-type: none"> <input type="checkbox"/> Yes <input type="checkbox"/> No 	Date progression was detected: Method used to detect progression (diagnostic testing):
	Please describe sites and symptoms associated with progression: Therapeutic plan: